National Library of Canada
Bibliothèque nationale du Canada

Canadian Theses Service
Service des thèses canadiennes

Ottawa, Canada
K1A 0N4

NOTICE

The quality of this microform is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us an inferior photocopy.

Reproduction in full or in part of this microform is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30, and subsequent amendments.

AVIS

La qualité de cette microforme dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de qualité inférieure.

La reproduction, même partielle, de cette microforme est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30, et ses amendements subséquents.

NL-339 (r. 86/04) c

Canada

THE UNIVERSITY OF ALBERTA

FINITELY GENERATED SOLUBLE GROUPS WITH IDENTITIES ON SUBGROUPS

BY

JAMES F. SEMPLE

A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES AND RESEARCH

IN PARTIAL FULFILLMENT OF THE REQUIREMENT FOR THE DEGREE

OF MASTER OF SCIENCE

DEPARTMENT OF MATHEMATICS

EDMONTON, ALBERTA

SPRING, 1989

National Library of Canada
Bibliothèque nationale du Canada

Canadian Theses Service
Service des thèses canadiennes

Ottawa, Canada
K1A 0N4

The author has granted an irrevocable non-exclusive licence allowing the National Library of Canada to reproduce, loan, distribute or sell copies of his/her thesis by any means and in any form or format, making this thesis available to interested persons.

L'auteur a accordé une licence irrévocable et non exclusive permettant à la Bibliothèque nationale du Canada de reproduire, prêter, distribuer ou vendre des copies de sa thèse de quelque manière et sous quelque forme que ce soit pour mettre des exemplaires de cette thèse à la disposition des personnes intéressées.

The author retains ownership of the copyright in his/her thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without his/her permission.

L'auteur conserve la propriété du droit d'auteur qui protège sa thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

ISBN 0-315-52816-8

Canada

THE UNIVERSITY OF ALBERTA

RELEASE FORM

NAME OF AUTHOR: JAMES F. SEMPLE

TITLE OF THESIS: FINITELY GENERATED SOLUBLE GROUPS WITH IDENTITIES ON SUBGROUPS

DEGREE FOR WHICH THESIS WAS PRESENTED: MASTER OF SCIENCE

YEAR THIS DEGREE GRANTED: 1989

Permission is hereby granted to THE UNIVERSITY OF ALBERTA LIBRARY to reproduce single copies of the thesis and to lend or sell such copies for private, scholarly or scientific research purposes only.

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.

James F. Semple

SIGNED

PERMANENT ADDRESS:

1897 Prince of Wales Drive,
Ottawa, Ontario
K2C 3J7

Date: 25 April 1989

THE UNIVERSITY OF ALBERTA

FACULTY OF GRADUATE STUDIES AND RESEARCH

The undersigned certify that they have read, and recommend to the Faculty of Graduate Studies and Research, for acceptance, a thesis entitled FINITELY GENERATED SOLUBLE GROUPS WITH IDENTITIES ON SUBGROUPS submitted by James F. Semple in partial fulfilment of the requirements for the degree of Master of Science.

Supervisor

Date: April 25, 1989

ABSTRACT

We call a group G an SP-group if there exist an integer $n > 1$ and distinct words u and v in the semigroup $\langle X_1, \ldots, X_n \mid X_i^2 = X_i \rangle$ such that the complexes $u(H_1, \ldots, H_n)$ and $v(H_1, \ldots, H_n)$ are equal; this may then be thought of as an identity on the subgroups of G. It is shown here that a finitely generated soluble SP-group is nilpotent-by-finite, and that any SP-group is either of restricted width or satisfies an identity in three variables.

ACKNOWLEDGEMENT

I must thank my supervisor, Professor Rhemtulla, who has demonstrated remarkable patience and enthusiasm throughout the course of this venture; furthermore I thank David Riley, Michel Sadelain, Michael Voralia and numerous others who have contributed in making my stay in Edmonton a pleasant one.

Finally, I thank the University of Alberta for its generous financial support.

TABLE OF CONTENTS

PAGE

INTRODUCTION 1

CHAPTER ONE

Examples and preliminary results 3

CHAPTER TWO

Restricting the structure of SP-groups 11

CHAPTER THREE

SP-groups are nilpotent-by-finite 19

REFERENCES 24

NOTATION

The following notation is used in the text of this thesis:

$H \wr K$, the wreath product of H and K

$H \rtimes K$, the split extension of H by K

$G' = [G, G]$, the derived subgroup of G

$G^{(\alpha)}$, the α^{th} term of the derived series of G

$\zeta(G)$, the centre of G

$\zeta_i(G)$, the i^{th} term of the upper central series of G, $(\zeta_0(G) = 1)$

$\gamma_i(G)$, the i^{th} term of the lower central series of G, $(\gamma_1(G) = G)$

max, the maximal condition on subgroups

max-n, the maximal condition on normal subgroups

$C_G(H)$, the centralizer of H in G

$N_G(H)$, the normalizer of H in G

INTRODUCTION

We call a group G an SP-group, for subgroup-permutable group, if there is some non-trivial identity on the subgroups of G — one like $(H_1H_2)^2 = (H_2H_1)^2$ for all $H_1,\ H_2 \leq G$. Our interest is in classifying groups which have identities on their subgroups but, in the context of this thesis, restrict ourselves to the case that G is finitely generated and soluble.

If u and v are distinct words in the semigroup $\langle X_1, \ldots, X_n \mid X_i^2 = X_i \rangle$, we say $G \in SP(\{u\}, \{v\})$ if $u(H_1, \ldots, H_n) = v(H_1, \ldots, H_n)$ for all sequences of subgroups $H_1, \ldots, H_n \leq G$. The somewhat cumbersome notation is maintained in this text because this problem is easily generalized in the following manner: classify groups G for which there exist finite disjoint sets U and V of words $\subset \langle X_1, \ldots, X_n \mid X_i^2 = X_i \rangle$ such that $G \in SP(U, V)$. In this greater generality, $G \in SP(U, V)$ means that for every n-tuple of subgroups $(H_1, \ldots, H_n)$ there exist $u \in U$ and $v \in V$ such that the complexes $u(H_1, \ldots, H_n)$ and $v(H_1, \ldots, H_n)$ are equal.

It should be emphasized here that throughout, when we write of identities on subgroups we mean indeed that the *complexes* are equal, not the weaker condition that the subgroups generated by $u(H_1, \ldots, H_n)$ and $v(H_1, \ldots, H_n)$ are equal.

The foundations of study of groups of this nature are due to Philip Hall, who considered groups with elliptically-embedded subgroups; certain such groups have the more restrictive property that for some integer n, $(HK)^n = (KH)^n$ for any subgroups H and K. Finitely generated and soluble groups of this property have recently been studied, albeit in more generality, by A.H. Rhemtulla and J.S. Wilson, ([3], [6]) who found them to be finite-by-nilpotent.

In this thesis we study the variation that G satifies an identity involving more than two subgroups at a time: groups $G \in SP(\{u\},\{v\})$ for which u and v are words in at least three letters.

In Chapter One, we reaffirm the notions introduced here, and prove some preliminary results; in particular it turns out that any group satisfying a subgroup identity satisfies one in at most three variables, and we exhibit such an identity on subgroups of the infinite dihedral group. Unfortunately D_∞ is not finite-by-nilpotent, so we aim to show that SP-groups have the weaker property of being nilpotent-by-finite.

The second chapter deals with a pair of key lemmas which are used to eliminate certain classes of groups, and therefore allow us a better picture of the structure and rank of SP-groups. These lemmas are then employed in the final chapter, which consists primarily of a proof by induction on its solubility length that our SP-group is nilpotent-by-finite.

CHAPTER ONE

EXAMPLES AND PRELIMINARY RESULTS

In this chapter, we give an example of an SP-group which is not finite-by-nilpotent, and we prove a number of preliminary results which are applicable to general groups with identities on their subgroups. We begin, however, by formalizing the concepts previously introduced.

[1.1] Definition: A subgroup H of a group G is said to be *elliptically embedded* in G if, for every $K \leq G$ there exists an integer n such that $\langle H, K \rangle = (HK)^n$. Groups all of whose subgroups are elliptically embedded are said to be of *restricted width*.

Examples of groups of restricted width are the so-called *quasi-hamiltonian groups*, that is groups G with the property that if $H, K \leq G$ then $HK = KH$. These quasi-hamiltonian groups are, in fact, examples of groups of restricted width which have the further property that there exists an integer n such that $(KH)^n = (HK)^n$ for all pairs H and $K \leq G$. It is this further property that we care to generalize, in the following manner:

[1.2] Definition: Let G be a group, then we say $G \in SP(\{u\}, \{v\})$ for words $u = u(X_1, X_2, \ldots, X_n) = X_{j_1} \cdots X_{j_s}$ and $v = v(X_1, X_2, \ldots, X_n) = X_{i_1} \cdots X_{i_r}$ if $u \neq v$ but the complexes $u(H_1, \ldots, H_n)$ and $v(H_1, \ldots, H_n)$ are equal for all sequences of subgroups $(H_1, \ldots, H_n)$; to guarantee that u and v are essentially different we further insist that $i_\ell \neq i_{\ell+1}$ and $j_\ell \neq j_{\ell+1}$ throughout. We refer to G

as an SP-group if there exist u and v with $G \in SP(\{u\},\{v\})$.

Therefore, particular examples of SP-groups are quasi-hamiltonian groups and groups of restricted width which have bounded ellipticity. It is our aim in this thesis to show that if G is a finitely-generated and soluble SP-group then G is nilpotent-by-finite; we begin with the following

[1.3] Lemma. *Let G be a group, and suppose there exists an integer n such that $G \in SP(\{u\},\{v\})$ for some $u = u(X_1, X_2, \ldots, X_n)$ and $v = v(X_1, X_2, \ldots, X_n)$, $u \neq v$. If n is the least such integer, then $n = 2$ or 3.*

Proof: Assume $n > 3$, and define

$$u_i(X_1, \ldots, X_{i-1}, X_{i+1}, \ldots, X_n) = u(X_1, \ldots, X_{i-1}, 1, X_{i+1}, \ldots, X_n).$$

Write each u_i in canonical form, replacing X_j^2 by X_j throughout.

By the minimality of n, we have that u_i and v_i are identically equal for all i; we claim this implies $u = v$ or, equivalently, that there is a unique word $w = w(X_1, X_2, \ldots, X_n)$ such that $w_i = u_i$ for $i = 1 \ldots n$

$$\begin{aligned}
\text{Let } u_1 &= X_{m_{1,1}} X_{m_{1,2}} \cdots X_{m_{1,\ell(1)}} \\
u_2 &= X_{m_{2,1}} X_{m_{2,2}} \cdots X_{m_{2,\ell(2)}} \\
u_3 &= X_{m_{3,1}} X_{m_{3,2}} \cdots X_{m_{3,\ell(3)}} \\
u_4 &= X_{m_{4,1}} X_{m_{4,2}} \cdots X_{m_{4,\ell(4)}} \quad \text{et cetera,}
\end{aligned}$$

and let $w = X_{a_1} X_{a_2} \ldots X_{a_k}$ have the property $w_i = u_i$ for all i.

We give the following algorithm to reconstruct w from the given set $\{w_i\}$:

$$u_1 \Longrightarrow a_1 = 1 \text{ or } a_1 = m_{1,1}$$

$$u_2 \Longrightarrow a_1 = 2 \text{ or } a_1 = m_{2,1}$$

$$u_3 \Longrightarrow a_1 = 3 \text{ or } a_1 = m_{3,1}$$

and these three conditions determine a_1 uniquely. Consider now u_{a_1}, and we see that $a_2 = m_{a_1,1}$. Henceforth, we proceed iteratively: to determine a_i given a_{i-1} and a_{i-2}, consider two rows other than $u_{a_{i-1}}$ and $u_{a_{i-2}}$, say u_r and u_s. Let X_{r_i} and X_{s_i}, respectively, be the first terms of u_r and u_s whose position in w has not been determined. Then $a_i = (r_i \text{ or } r)$ and $(s_i \text{ or } s)$; this defines a_i uniquely unless $s = r_i$ and $r = s_i$. In this case, check the first term of u_{a_i} following the term $X_{a_{i-1}}$. It is clear that these three conditions determine a_i uniquely, and hence that the pre-image, w, of the "projections" is unique. Therefore $u = v$, contrary to the requirement that $u \neq v$ if $G \in SP(\{u\}, \{v\})$.

We illustrate the algorithm as follows:

$$\text{Let } u_1 = X_2X_3X_4$$

$$u_2 = X_1X_3X_4$$

$$u_3 = X_1X_2X_1X_4$$

$$u_4 = X_1X_2X_1X_3$$

If $u(X_1, X_2, X_3, X_4) = X_{a_1}X_{a_2}\ldots X_{a_\ell}$, then u_1 tells us $a_1 \in \{1, 2\}$ (as does u_2), while u_3 determines that $a_1 = 1$.

Consider now u_1, and we see that $a_2 = 2$. Now we look at both u_3 and u_4

to see that a_3 is in both $\{3,1\}$ and $\{4,1\}$, that is we must have $a_3 = 1$. Finally we look again at u_3 and u_4 to see that a_4 is in both $\{3,4\}$ and $\{4,3\}$. This is inconclusive, so we check u_2 for the first entry not accounted for — in this case X_3. We continue in this manner to determine the unique word $u = X_1X_2X_1X_3X_4$ whose "projection by i" is, in each case, u_i.

[1.4]Remark: If G is finitely-generated and soluble, and $G \in SP(\{u\},\{v\})$ for some words u and v in two variables, then G is of restricted width. The result of Rhemtulla [3] implies, therefore, that G is finite-by-nilpotent. We apply a result due to P. Hall ([8], part I, p.117) which states that if for a group G there exists an integer i such that $\gamma_i(G)$ is finite, then there is an integer j such that $|G : \zeta_j(G)|$ is finite. We clearly have an ascending central series to $\zeta_j(G)$, so G is nilpotent-by-finite.

Therefore, we have in particular that finite-by-nilpotent groups are nilpotent-by-finite, and it follows that so too are finitely generated soluble SP-groups in the case that our words are in two variables. Thus, we need here address only the case that u and v are words in three variables.

[1.5]Lemma. *Let $G \in SP(\{u\},\{v\})$, with $u(X_1,X_2,X_3) = X_{i_1}X_{i_2}\cdots X_{i_m}$ and $v(X_1,X_2,X_3) = X_{j_1}X_{j_2}\cdots X_{j_n}$. Then we may assume that each of X_1, X_2 and X_3 appear the same number of times in the expressions for u and v.*

Proof: Without loss, assume that X_1 appears more often in $u(X_1,X_2,X_3)$ than in $v(X_1,X_2,X_3)$. Then $u(X_1,X_2,X_2) \neq v(X_1,X_2,X_2)$, (replacing X_2^2 by X_2 throughout,) so if we let $u'(X_1,X_2) = u(X_1,X_2,X_2)$, and similarly $v'(X_1,X_2) =$

$v(X_1, X_2, X_2)$ then $G \in SP(\{u'\}, \{v'\})$. Therefore G is of restricted width, so by the previous remark G is nilpotent-by-finite.

Corollary: If G is an SP-group for words $u(X_1, X_2, X_3) = X_{i_1} X_{i_2} \cdots X_{i_m}$ and $v(X_1, X_2, X_3) = X_{j_1} X_{j_2} \cdots X_{j_n}$ then G is nilpotent-by-finite or $m = n$.

A similarly simple but useful result is

[1.6]Lemma. *Let $G \in SP(\{u\}, \{v\})$ with $u(X_1, X_2, X_3) = X_{i_1} X_{i_2} \cdots X_{i_m}$ and $v(X_1, X_2, X_3) = X_{j_1} X_{j_2} \cdots X_{j_n}$. Then we may assume that $i_1 = j_1$ and $i_n = j_n$.*

Proof: Consider the case that $i_1 \neq j_1$. Without loss of generality, assume that $i_1 = 1$ and $j_1 = 2$. Once again if we let $u'(X_1, X_2) = u(X_1, X_2, X_2)$, and $v'(X_1, X_2) = v(X_1, X_2, X_2)$ and replace ${X_2}^2$ with X_2 as required, we find that $u'(X_1, X_2) \neq v'(X_1, X_2)$ and that $G \in SP(\{u'\}, \{v'\})$; hence, G is of restricted width, and is therefore nilpotent-by-finite.

We now prove the following result, which will be applied to prove that the infinite dihedral group is an SP-group:

[1.7]Lemma. *Let $G = D_\infty = \langle a, b \mid a^b = a^{-1}, b^2 = 1 \rangle$. If H is a subgroup of G which is not normal, then $H = \langle a^{r_1}, a^{s_1} b \rangle$ for some integers r_1 and s_1*

Proof: Let $R = \{r \in \mathbf{Z} \mid a^r \in H\}$, and $S = \{s \in \mathbf{Z} \mid a^s b \in H\}$. We have that $0 \in R$, so R is non-empty, and S is non-empty for otherwise $H \lhd G$.

Now $\{a^r \mid r \in R\} \leq H$, so there exists some $r_1 \in R$ with $\{a^r \mid r \in R\} = \langle a^{r_1} \rangle$. If $r_1 = 0$ then $H = \langle 1, a^{s_1} b \rangle$ and we are done. Otherwise, let s_1, $s_2 \in S$;

then $a^{s_1}ba^{s_2}b \in \langle a^{r_1} \rangle$, so

$$s_1 - s_2 \equiv 0 \,(mod\, r_1).$$

If, on the other hand, $s_1 \in \mathcal{S}$ and $s \equiv s_1 (mod\, r)$ then $s \in S$. Therefore given any $s_1 \in S$, S then equals $\{s \in \mathbf{Z} \mid s \equiv s_1 (mod\, r_1)\}$. From this we infer that

$$H = \langle a^{r_1}, a^{s_1}b \rangle = \langle a^{r_1} \rangle \rtimes \{1, a^{s_1}b\}.$$

This completes the proof of [1.7]

[1.8]Lemma. *There exist words u and v such that $D_\infty \in SP(\{u\}, \{v\})$.*

Proof: Let

$$u(X_1, X_2, X_3) = (X_1X_2)^2X_3(X_1X_2),$$
$$v(X_1, X_2, X_3) = (X_1X_2)X_3(X_1X_2)^2;$$

we claim $D_\infty \in SP(\{u\}, \{v\})$. Let $G = D_\infty$ and let H_1, H_2 and H_3 be any three subgroups of G. If any of the H_i is normal in G, then clearly $u(H_1, H_2, H_3) = v(H_1, H_2, H_3)$. Therefore we may assume no H_i is normal in G.

We must show that $u(H_1, H_2, H_3) = v(H_1, H_2, H_3)$ in the case that each $H_i = \langle a^{r_i} \rangle \rtimes \{1, a^{s_i}b\}$. We do this by showing first that we may assume each H_i has order two. Denote $A_i = \langle a^{r_i} \rangle$, $B_i = \{1, a^{s_i}b\}$, then $H_i = A_iB_i$. Now each $A_i \lhd G$ and, if any H_j is of infinite order we may replace G by G/A_j and each H_i by H_iA_j/A_j. It suffices therefore to show that $u(B_1, B_2, B_3) = v(B_1, B_2, B_3)$ — that is, we need only to prove the result for assuming our subgroups are of order 2.

We show here that $u(B_1, B_2, B_3) \subseteq v(B_1, B_2, B_3)$, for the proof of the converse is similar:

Let U and V denote $u(B_1, B_2, B_3)$ and $v(B_1, B_2, B_3)$, respectively. Consider a word $w = b_1b_2b_1'b_2'b_3b_1''b_2'' \in U$, with each $b_j^{(n)} \in B_j$, for $n \in \{0, 1, 2\}$. If $b_3 = 1$

then $w \in V$, so we may assume that $b_3 = a^{s_3}b$; moreover, $b_1 b_2 b_1' b_2'$ is necessarily in $B_1 B_2$ except in the cases:

$$(i) \quad b_1 = 1,\ b_2 = a^{s_2}b,\ b_1' = a^{s_1}b,\ b_2' = 1$$

$$(ii) \quad b_1 = 1,\ b_2 = a^{s_2}b,\ b_1' = a^{s_1}b,\ b_2' = a^{s_2}b$$

$$(iii) \quad b_1 = a^{s_1}b,\ b_2 = a^{s_2}b,\ b_1' = a^{s_1}b,\ b_2' = 1$$

$$(iv) \quad b_1 = a^{s_1}b,\ b_2 = a^{s_2}b,\ b_1' = a^{s_1}b,\ b_2' = a^{s_2}b$$

We shall make use of the following identity:

$$a^{s_2}ba^{s_1}ba^{s_3}b \ = \ a^{s_3}ba^{s_1}ba^{s_2}b.$$

Therefore, in the case

$$\begin{aligned}(i)\ b_1 b_2 b_1' b_2' b_3 B_1 B_2 &= a^{s_2}ba^{s_1}ba^{s_3}bB_1B_2 \\ &= a^{s_3}ba^{s_1}ba^{s_2}bB_1B_2 \\ &\subset B_3(B_1B_2)^2.\end{aligned}$$

$$\begin{aligned}(ii)\ b_1 b_2 b_1' b_2' b_3 B_1 B_2 &= a^{s_2}ba^{s_1}ba^{s_2}ba^{s_3}bB_1B_2 \\ &= a^{s_2}ba^{s_3}ba^{s_2}ba^{s_1}bB_1B_2 \\ &\subset a^{s_2}ba^{s_3}ba^{s_2}bB_1B_2 \\ &\subset B_2B_3(B_1B_2)^2.\end{aligned}$$

$$\begin{aligned}(iii)\ b_1 b_2 b_1' b_2' b_3 B_1 B_2 &= a^{s_1}ba^{s_2}ba^{s_1}ba^{s_3}bB_1B_2 \\ &\subset a^{s_1}bB_3(B_1B_2)^2, \text{ using } (i) \\ &\subset B_1B_3(B_1B_2)^2.\end{aligned}$$

$$(iv)\ b_1b_2b_1'b_2'b_3B_1B_2 = a^{s_1}bB_2B_3(B_1B_2)^2,\ \text{using}(ii)$$

$$\subset B_1B_2B_3(B_1B_2)^2.$$

In each case, we have $b_1b_2b_1'b_2'b_3B_1B_2 \subset B_1B_2B_3(B_1B_2)^2$, that is, $U \subseteq V$. The proof of the opposite inclusion is similar.Therefore $u(H_1, H_2, H_3) = v(H_1, H_2, H_3)$ for all subgroups H_1, H_2, and H_3.

Therefore, there exist words u and v such that $D_\infty \in SP(\{u\}, \{v\})$. One will note, however, that D_∞ is nilpotent-by-finite, but is not finite-by-nilpotent; hence, it is an example of a group which satisfies an identity on its subgroups, but for which not all subgroups are elliptically embedded.

CHAPTER TWO

RESTRICTING THE STRUCTURE OF *SP*-GROUPS

In this chapter we establish a pair of key lemmas to consider the structure of SP-groups. In Chapter Three we shall make inferences about the rank of G from Lemma [2.2],which states that our SP-groups are minimax. To prove this lemma we make use of the following theorem due to P.H. Kropholler [2].

[2.1]Theorem. *If G is a finitely generated soluble group, and if for no prime p does G have a section isomorphic to $C_p \wr C_\infty$, then G is minimax.*

[2.2]Lemma. *If G is a finitely generated soluble group, and $G \in SP(\{u\},\{v\})$ for some $u \neq v$ then G is minimax.*

Proof: By the Kropholler's theorem, it is sufficient to show that G has no section isomorphic to $C_p \wr C_\infty$; however, because $G \in SP(\{u\},\{v\})$ is a quotient- and subgroup-closed property it suffices to show that for distinct words u and v, $C_p \wr C_\infty \notin SP(\{u\},\{v\})$.

Assume that $C_p \wr C_\infty \in SP(\{u\},\{v\})$ for $u(X_1,X_2,X_3) = X_{i_1}X_{i_2}\cdots X_{i_m}$ and $v(X_1,X_2,X_3) = X_{j_1}X_{j_2}\cdots X_{j_m}$. With no loss in generality we may further assume that $i_1 = j_1 = 1$.

We claim first that $C_p \wr C_\infty$ is isomorphic to $\mathbf{F_p}\langle x\rangle \rtimes \langle t\rangle$, where x and t are of infinite order and the action of t on $a(x) \in \mathbf{F_p}\langle x\rangle$ is multiplication by x.

We begin by showing that the base group B of $C_p \wr C_\infty$ is isomorphic to $\mathbf{F_p}\langle x\rangle$. For each $r \in \{0,\ldots,p-1\}$ and each $s \in \mathbf{Z}$, define a permutation $\theta_{r,s}$ of

$C_p \times C_\infty$ in the following manner: for $(a^\ell, \mu^m) \in C_p \times C_\infty$,

$$\theta_{r,s}(a^\ell, \mu^m) = \begin{cases} (a^{\ell+s}, \mu^m) & \text{if } m = s; \\ (a^\ell, \mu^m) & \text{if } m \neq s; \end{cases}$$

The set of all $\theta_{r,s}$ generates the base group, which is clearly abelian. Let

$$B = \langle \theta_{r,s} \mid r \in \{0, \ldots p-1\}, s \in \mathbf{Z} \rangle.$$

Define $\psi: B \longrightarrow \mathbf{F_p}\langle x \rangle$ by $\psi(\theta_{r_1,s_1} \cdots \theta_{r_n,s_n}) = \sum_{i=1}^{n} r_i x^{s_i}$. It is immediate that ψ is onto and homomorphic, while the facts that B is abelian and that $\theta_{u,s} \cdot \theta_{v,s} = \theta_{u+v,s}$ together yield that ψ is well-defined and one-to-one. Therefore $B \cong \mathbf{F_p}\langle x \rangle$.

For integers v define $\sigma_v \in C_p \wr C_\infty$ by $\sigma_v(a^\ell, \mu^m) = (a^\ell, \mu^{m-v})$. Then $\langle \sigma_v \mid v \in \mathbf{Z} \rangle = \langle \sigma_1 \rangle$ and, by definition of the wreath product,

$$C_p \wr C_\infty \cong B \rtimes \langle \sigma_1 \rangle.$$

Note that $(\theta_{r,s})^{\sigma_v} = \theta_{r,s+v}$. If we extend ψ to ψ': $C_p \wr C_\infty \rightarrow F_p\langle x \rangle \rtimes \langle t \rangle$ by $\psi'(\sigma_r) = t^r$, then we have our desired isomorphism from $C_p \wr C_\infty$ to $\mathbf{F_p}\langle x \rangle \rtimes \langle t \rangle$.

For $a_i(x) \in \mathbf{F_p}\langle x \rangle$ and $c_i = (a_i(x)(1 - x^{m_i}), t^{-m_i}) \in C_p \wr C_\infty$, denote

$$C_i = \langle c_i \rangle = \left\{ (a_i(x)(1 - x^{k_i m_i}), t^{-k_i m_i}) \mid k_i \in \mathbf{Z} \right\}$$

Consider a general term, w, of $C_{i_1} C_{i_2} \cdots C_{i_\ell}$:

$$w = \big(a_{i_1}(x)(1-x^{k_1 m_{i_1}}), t^{-k_1 m_{i_1}}\big)\big(a_{i_2}(x)(1-x^{k_2 m_{i_2}}), t^{-k_2 m_{i_2}}\big)\cdots$$

$$\cdots\big(a_{i_\ell}(x)(1-x^{k_\ell m_{i_\ell}}), t^{-k_\ell m_{i_\ell}}\big)$$

$$= \big(a_{i_1}(x)(1-x^{k_1 m_{i_1}}) + x^{k_1 m_{i_1}} a_{i_2}(x)(1-x^{k_2 m_{i_2}}), t^{-k_1 m_{i_1} - k_2 m_{i_2}}\big)\cdots$$

$$\cdots\big(a_{i_\ell}(x)(1-x^{k_\ell m_{i_\ell}}), t^{-k_\ell m_{i_\ell}}\big)$$

$$\vdots$$

$$= \big(a_{i_1}(x)(1-x^{k_1 m_{i_1}}) + \ldots$$

$$\ldots + x^{k_1 m_{i_1} + \ldots + k_{\ell-1} m_{i_{\ell-1}}} a_{i_\ell}(x)(1-x^{k_\ell m_{i_\ell}}), t^{-k_1 m_1 - \ldots - k_\ell m_{i_\ell}}\big)$$

Let us denote $\sum_{j=1}^{s} k_j m_{i_j}$ by $d(s)$, (with $d(0)=0$), and, for $\ell \in \{1,2,3\}$ define $I_\ell = \{r \in \mathbf{Z} \mid a_{i_r} = a_\ell\}$. Consider now the component of w in the base group; it equals

$$a_1(x)\sum_{j\in I_1}\big(x^{d(j-1)} - x^{d(j)}\big) + a_2(x)\sum_{j\in I_2}\big(x^{d(j-1)} - x^{d(j)}\big) + a_3(x)\sum_{j\in I_3}\big(x^{d(j-1)} - x^{d(j)}\big).$$

Choose primes p_1, p_2, p_3, p_4 with $p_4 > p_3 > p_2 > p_1 > m$, and let $m_1 = p_2p_3p_4$, $m_2 = p_1p_3p_4$, $m_3 = p_1p_2p_4$ and $a_1(x) = 1$, $a_2(x) = x^{p_1+1}$, $a_3(x) = x^{p_2+1}$. Take $k_1 = \ldots = k_m = 1$, and let k'_1, k'_2, $\ldots$ k'_m be such that

$$\sum_{i=1,2,3} a_i(x)\sum_{j\in I_i}\big(x^{d(j-1)} - x^{d(j)}\big) = \sum_{i=1,2,3} a_i(x)\sum_{j\in I'_i}\big(x^{d'(j-1)} - x^{d'(j)}\big)$$

where the d', I'_i and k' of the right-hand side are analogous to the d, I_i and k of the left.

We aim first to show that there are $2m$ powers of x appearing in the left sum; certainly there are no more than $2m$ terms.

If there is to be cancellation, we must have a term x^r which is expressed as both $a_{\ell_i}(x)x^{d(s)}$ and $a_{\ell_j}(x)x^{d(t)}$. Let $a_{\ell_i}(x) = x^{b_i}$ and $a_{\ell_j}(x) = x^{b_j}$, where $b_i, b_j \in \{1, p_1+1, p_2+1\}$. Now $d(t) \equiv d(s) \equiv 0 \,(mod\, p_4)$ implies $b_i \equiv b_j \,(mod\, p_4)$. The inequality $p_4 > p_3 > p_2 > p_1 > 1$ implies further that $a_{\ell_i}(x) = a_{\ell_j}(x)$.

Moreover $d : \{0, \ldots, m\} \to \mathbf{N}$ is monotonically increasing, so $d(s) = d(t)$ implies $s = t$. Therefore a term $a_{\ell_i}(x)x^{d(s)}$ can be cancelled only by subtracting $a_{\ell_i}(x)x^{d(s)}$, but the assumption $i_t \not\equiv i_{t+1}$ rules out this possibility. Therefore, there are precisely $2m$ powers of x appearing in the sum on the left.

Note that p_4 divides each of the powers $d(j)$ and that the $a_i(x)$ have powers incongruent mod p_4. It follows that, for each i,

$$\sum_{j \in I_i} \left(x^{d(j-1)} - x^{d(j)}\right) = \sum_{j \in I'_i} \left(x^{d'(j-1)} - x^{d'(j)}\right)$$

We wish to show now that we must have $k'_1 = k'_2 = \ldots = k'_n = 1$ and $i_\ell = j_\ell$ for $\ell = 1 \ldots n$. It suffices to show that $d(\ell) = d'(\ell)$ for $\ell = 1 \ldots m$.

Assume, by way of contradiction, that $k'_1 \neq 1$; however $-a_1(x)x^{d'(1)}$ appears in the right-hand expression, so for some $t > 1$ we have $d'(1) = d(t)$, ie.,

$$m_1 k'_1 = m_1 + m_{i_2} + \ldots + m_{i_t}.$$

Now $i_2 \neq 1$, and p_{i_2} divides m_{i_ℓ} if $i_\ell \neq i_2$. If β is the number of $b \leq t$ for which $m_{i_\ell} = m_{i_2}$, then we have

$$m_{i_2}\beta \equiv 0 \,(mod\, p_2)$$

However $gcd(m_{i_2}, p_{i_2}) = 1$ and $p_{i_2} > \beta \geq 0$ imply $\beta = 0$. Hence $m_1 k'_1 = m_1$. This implies that $k'_1 = 1$ and that $d(1) = d'(1)$. Conversely if $d(1) = d'(1)$ and

$d(1) = d'(t)$ then, because there is no cancellation on the left-hand side and d is monotonically increasing, we must have $t = 1$.

We proceed by induction as follows:

Assume for $r \in \{1, \ldots, s\}$ that

$$d'(r) = d(t) \iff r = t$$

Now $d'(s+1) = d(t)$ for some $t \geq s+1$.

$$d'(s+1) = d'(s) + m_{j_{s+1}} k'_{s+1} = d(s) + m_{j_{s+1}} k'_{s+1}$$

Therefore $m_{j_{s+1}} k'_{s+1} = (m_{i_{s+1}} + m_{i_{s+2}} + \ldots + m_{i_t})$. If $j_{s+1} = i_{s+1}$ then $p_{i_{s+2}}$ divides $m_{j_{s+1}}$. Our approach is as with the case $k'_1 = 1$:

$$p_{i_{s+2}} \text{ divides } \underbrace{(1 + 1 + \ldots + 1)}_{<t} m_{i_{s+2}} \Rightarrow t = s+1$$

Therefore $k'_{s+1} = 1$ and $j_{s+1} = i_{s+1}$

If on the other hand $j_{s+1} \neq i_{s+1}$, we proceed similarly:

$p_{i_{s+1}}$ divides $m_{i_{s+1}}(1 + \ldots + 1)$ so $m_{j_{s+1}} k'_{s+1} = 0$, and therefore $k'_{s+1} = 0$. In this case

$$\left\langle (a_{j_{s+1}}(x)(1 - x^{k'_{s+1} m_{j_{s+1}}}), t^{-k_{s+1} m_{j_{s+1}}}) \right\rangle = 1$$

and we have fewer than $2m$ powers of x appearing on the right-hand side.

Therefore $m_{j_{s+1}} = m_{i_{s+1}}$. By induction, we have $j_s = i_s$ for $s = 1, \ldots, n$. This implies that $u(X_1, X_2, X_3) = v(X_1, X_2, X_3)$, as required. We conclude that for all u and v we have $C_p \wr C_\infty \notin SP(\{u\}, \{v\})$.

We will make use of the following lemma, which is a slight modification of Lemma 4 of [4] and its corollary:

[2.3]Lemma. *Let α be a non-zero algebraic number such that for some fixed β the equation*

$$\alpha^{\lambda_1} \pm \alpha^{\lambda_2} \pm \ldots \pm \alpha^{\lambda_m} = \beta$$

has solutions in rational integers $\lambda_1, \ldots, \lambda_m$ together with the following further property: For a given solution, if Λ is the set of those λ_i for which α^{λ_i} does not appear as many times with positive as negative coefficient, then $\lambda = max\{\lambda_i \in \Lambda\}$ takes on arbitrarily large positive values. If α and all its non-zero integral powers satisfy these hypotheses, then α is a root of unity.

Proof: As in ([4], Lemma 4) we have that for every absolute value $|\ |_v$ of $\mathbf{Q}(\alpha)$,

$$|\alpha|_v \leq f_v(m) = \begin{cases} m, & \text{if } v \text{ archimedean;} \\ max\{|r|_v^{-1}, r = 1, \ldots, m\}, & v \text{ non-archimedean.} \end{cases}$$

Now $|\alpha^k|_v \leq f_v(m)$ for all non-zero integers k, so taking k^{th} roots and letting $k \to \infty$ we find that $|\alpha|_v \leq 1$ for every absolute value of $\mathbf{Q}(\alpha)$; however taking $k \to -\infty$ we find that $|\alpha|_v \geq 1$ for every absolute value of $\mathbf{Q}(\alpha)$. It is a classical result of algebraic number theory that if α is algebraic and is not a root of unity then there exists an absolute value such that $|\alpha|_v \neq 1$. (See for example [1], p.109) Therefore α is a root of unity, as desired.

[2.4]Lemma. *Let α be algebraic over $\mathbf{Q}$, and let $G_1 = \mathbf{Q}\langle\alpha\rangle \rtimes \langle t\rangle$ with the action of t on an $f(\alpha) \in \mathbf{Q}\langle\alpha\rangle$ being multiplication by α. If $G \leq G_1$ is a subgroup containing $\mathbf{Z}\langle\alpha\rangle \rtimes \langle t\rangle$, and $G \in SP(\{u\},\{v\})$ for some u and v, then some power of t centralizes G.*

Proof: The proof follows similar lines to the proof of [2.2]. We assume that $G \in SP(\{u\},\{v\})$ for some $u(X_1,X_2,X_3)= X_{i_1}X_{i_2}\cdots X_{i_n}$ and $v(X_1,X_2,X_3) = X_{j_1}X_{j_2}\cdots X_{j_n}$; without loss of generality we again assume that $i_1 = j_1 = 1$.

For $f_i(\alpha) \in \mathbf{Q}\langle\alpha\rangle$ and $c_i = \left(f_i(\alpha)(1-\alpha^{m_i}), t^{-m_i}\right) \in G$, denote

$$C_i = \langle c_i\rangle = \left\{ \left(f_i(\alpha)(1-\alpha^{k_i m_i}), t^{-k_i m_i}\right) \mid k_i \in \mathbf{Z} \right\}$$

We consider a general term, w, of $C_{i_1}C_{i_2}\ldots C_{i_\ell}$:

$$\begin{aligned} w &= \left(f_{i_1}(\alpha)(1-\alpha^{k_1 m_{i_1}}), t^{-k_1 m_{i_1}}\right)\left(f_{i_2}(\alpha)(1-\alpha^{k_2 m_{i_2}}), t^{-k_2 m_{i_2}}\right)\cdots \\ &\qquad \cdots \left(f_{i_\ell}(\alpha)(1-\alpha^{k_\ell m_{i_\ell}}), t^{-k_\ell m_{i_\ell}}\right) \\ &= \left(f_{i_1}(\alpha)(1-\alpha^{k_1 m_{i_1}}) + x^{k_1 m_{i_1}}\left(f_{i_2}(\alpha)(1-\alpha^{k_2 m_{i_2}}), t^{-k_1 m_{i_1} - k_2 m_{i_2}}\right)\cdots \right. \\ &\qquad \left. \cdots \left(f_{i_\ell}(\alpha)(1-\alpha^{k_\ell m_{i_\ell}}), t^{-k_\ell m_{i_\ell}}\right)\right. \\ &\quad \vdots \\ &= \left(f_{i_1}(\alpha)(1-\alpha^{k_1 m_{i_1}}) + \ldots \right. \\ &\qquad \left. \ldots + \alpha^{k_1 m_{i_1} + \ldots + k_{\ell-1} m_{i_{\ell-1}}} f_{i_\ell}(\alpha)(1-\alpha^{k_\ell m_{i_\ell}}), t^{-k_1 m_1 - \ldots - k_\ell m_{i_\ell}}\right) \end{aligned}$$

We maintain the notations for d, I_i, k, d', I_i' and k' from the proof of Lemma [2.2]. In partcular, $d(s) = \sum_{j=1}^{s} k_j m_{i_j}$, and the general form of the component of w in $Q(\alpha)$ is:

$$f_1(\alpha)\sum_{j\in I_1}\left(\alpha^{d(j-1)} - \alpha^{d(j)}\right) + f_2(\alpha)\sum_{j\in I_2}\left(\alpha^{d(j-1)} - \alpha^{d(j)}\right) + f_3(\alpha)\sum_{j\in I_3}\left(\alpha^{d(j-1)} - \alpha^{d(j)}\right).$$

We assume that for a choice of integers $k_1,\ldots k_n$ and elements $f_1(\alpha)$, $f_2(\alpha)$, $f_3(\alpha)$ of $\mathbf{Q}\langle\alpha\rangle$ there exist integers $k_1',\ldots k_n'$ such that

$$\sum_{i=1,2,3} f_i(\alpha)\sum_{j\in I_i}\left(\alpha^{d(j-1)} - \alpha^{d(j)}\right) = \sum_{i=1,2,3} f_i(\alpha)\sum_{j\in I_i'}\left(\alpha^{d'(j-1)} - \alpha^{d'(j)}\right)$$

For a given $k_1, \ldots k_n$ fix $k_1', \ldots k_n'$ and set

$$\varphi_k(x) = \sum_{i=1,2,3} f_i(x) \sum_{j \in I_i} \left(x^{d(j-1)} - x^{d(j)}\right) - \sum_{i=1,2,3} f_i(x) \sum_{j \in I_i'} \left(x^{d'(j-1)} - x^{d'(j)}\right)$$

We use the expression φ_k to remind ourselves that φ is dependent upon our choice of the k_i.

In our proof of Lemma [2.2] we showed that if we choose $k_1 = \ldots = k_n = 1$, and our f_i appropriately then the corresponding $\varphi_k(x)$ is identically zero if and only if $k_1' = \ldots = k_n' = 1$, viz. $u(X_1, X_2, X_3) = v(X_1, X_2, X_3)$; however inspection of the proof of [2.2] shows that we may relax our hypothesis somewhat:

Given primes $p_4 > p_3 > p_2 > p_1 > n$, choose n integers k_i each of the form $1 + b_i p_1 p_2 p_3 p_4$ for integers b_i.

G contains $\mathbf{Z}\langle\alpha\rangle \rtimes \langle t\rangle$, and in particular contains 1, α^{p_1+1}, and α^{p_2+1}; therefore, take $f_1(x) = 1$, $f_2(x) = x^{p_1+1}$, $f_3(x) = x^{p_2+1}$. If the corresponding $\varphi_k(x)$ is identically zero then $u(X_1, X_2, X_3) = v(X_1, X_2, X_3)$.

In particular if we choose any integers b_i with $1 < b_1 < \ldots < b_n$, and set $k_i = 1 + b_i p_1 p_2 p_3 p_4$, then we cannot have $\varphi_k(x)$ identically zero. Every $\varphi_k(\alpha) = 0$, and clearly as we take arbitrarily large values for the b_i we find the powers appearing in $\varphi_k(x)$ to grow arbitrarily large. Therefore, α meets the requirements of the hypotheses of Lemma [2.3].

Furthermore, for every non-zero ℓ we find that α^ℓ satisfies the hypotheses of this lemma: if $G \leq \mathbf{Q}\langle\alpha\rangle \rtimes \langle t\rangle$ is an SP-group, then so too will be $G \bigcap \mathbf{Q}\langle\alpha^\ell\rangle \rtimes \langle t^\ell\rangle$. If we now mimic the proof that α satisfies the hypotheses of the aforementioned lemma, we find that α^ℓ does, too.

Therefore, by Lemma [2.3] α is a root of unity. As the action of t on $\mathbf{Q}\langle\alpha\rangle$ is multiplication by α, it follows that a power of t centralizes G.

CHAPTER THREE

SP-GROUPS ARE NILPOTENT-BY-FINITE

In this chapter we prove by induction on its solubility length that a finitely-generated SP-group is nilpotent-by-finite. We begin with a number of lemmas which will be employed in the course of the main proof.

[3.1]Lemma. *Let H be a torsion-free nilpotent group. Then $\zeta(H)$ is isolated. (i.e., $H/\zeta(H)$ is torsion-free.)*

Proof: Assume $g^p \in \zeta(H)$, let $\zeta_1 \leq \zeta_2 \leq \ldots \leq H$ be the upper central series of H, and let i be the least integer such that $[g,\zeta_i] \neq 1$. Take $x \in \zeta_i$ so $[g,x] \neq 1$. Then

$$1 = [g^p, x] = \left([g,x]^{g^{p-1}} \cdots [g,x]\right)$$

but

$$[g,x] \in \zeta_{i-1}, \text{ so } [g,x]^g = [g,x].$$

Now $1 = [g,x]^p$ and the fact that H is torsion-free together imply that $[g,x] = 1$. Therefore $g \in \zeta(H)$.

An immediate corollary is that if H is torsion-free, then every central factor of H is torsion-free.

[3.2]Lemma. *Let H be a torsion-free finitely-generated nilpotent group. Then H has a central series with infinite-cyclic factors.*

Proof: Consider the upper central series for H. By lemma [3.1], each factor is torsion-free; moreover H satisfies *max*, so the upper central factors are in fact

finitely-generated and torsion-free abelian. We refine this series to get one with infinite cyclic factors.

[3.3]Lemma. *If G is a polycyclic group, then G is poly(C_∞)-by-finite.*

Proof: Let $1 = G_0 \lhd G_1 \lhd \cdots \lhd G_r = G$ be a finite series with cyclic factors. The proof is by induction on r, with the case $r = 1$ holding true trivially.

Assume that K is a poly(C_∞) normal subgroup of G_{r-1} with G_{r-1}/K finite. Let $K^\circ = \bigcap_{g \in G} K^g$, the core of K in G. Then G_{r-1}/K° is finitely generated, torsion and soluble, so it is finite.

If G_r/G_{r-1} is finite, then the fact that K is poly(C_∞) implies that K° is poly(C_∞) and certainly $K^\circ \lhd G_r$, so K° is a poly(C_∞) group of finite index in G. If, alternatively, $G_r/G_{r-1} \cong C_\infty$, write $G/G_{r-1} = \langle xG_{r-1} \rangle$.

Now G_{r-1}/K° is finite, and so too must $Aut(G_{r-1}/K^\circ)$ be finite. Each element of $\{x, x^2, \dots\}$ induces an automorphism of G_{r-1}/K°, so x^m centralizes G_{r-1}/K° for some positive m. Let $L = \langle x^m, K^\circ \rangle$, then $L \lhd G$. Considering the Hirsch lengths we see that G/L is finite, and that L/K° is infinite cyclic. Therefore G contains a poly(C_∞) group of finite index, as required.

[3.4]Lemma. *Let A be a torsion-free abelian group of finite rank, and consider an SP-group $G = A \rtimes \langle t \rangle$. Then $\zeta(\langle A, t^\ell \rangle) \bigcap A \neq 1$ for some $\ell > 0$.*

Proof: Extend the action of $\langle t \rangle$ on A to $V = A \otimes_{\mathbf{Z}} \mathbf{Q}$. We write the action of t on $A \otimes_{\mathbf{Z}} \mathbf{Q}$ as multiplication, that is given any $f(t) \in \mathbf{Q}\langle t \rangle$, the map $a \otimes 1 \to (a \otimes 1) \cdot f(t)$ is an endomorphism of $A \otimes_{\mathbf{Z}} \mathbf{Q}$. Therefore $A \otimes_{\mathbf{Z}} \mathbf{Q}$ is a $\mathbf{Q}\langle t \rangle$-module.

Let $V_1 = A_1 \otimes_{\mathbf{Z}} \mathbf{Q}$ be an irreducible $\mathbf{Q}\langle t \rangle$-submodule of V. By Schur's Lemma,

$\Gamma = \mathrm{End}_{\mathbf{Q}\langle t\rangle} V_1$ is a division ring, finite dimensional over $\mathbf{Q}$. Considered as elements of $\mathrm{End}_{\mathbf{Q}} V_1$, $\langle t\rangle$ spans $\mathrm{End}_{\mathbf{Q}\langle t\rangle} V_1$, so Γ is commutative. It is therefore an algebraic number field.

V_1 is one-dimensional as a Γ-space as it is irreducible, so we may identify conjugation of V_1 by $\langle t\rangle$ with multiplication by an algebraic number, τ. We may now think of A_1 as a subgroup of $\mathbf{Q}(\tau)$ under addition.

Moreover A_1 is an SP-group, so by Lemma [2.4] there is an integer ℓ such that t^ℓ centralizes A_1. Therefore $A_1 \subset \zeta(\langle A, t^\ell\rangle) \bigcap A$. Hence, $\zeta(\langle A, t^\ell\rangle) \bigcap A \neq 1$, as was required.

[3.5]Theorem. *Let G be a finitely-generated soluble group, $G \in SP(\{u\},\{v\})$ for some u and v, then G is nilpotent-by-finite.*

Proof: We proceed by induction on the solubility length of G, with the result holding trivially for abelian groups. Assume that G has solubility length ℓ, and let $A = G^{(\ell-1)}$. Now G/A is finitely-generated and soluble, and $G/A \in SP(\{u\},\{v\})$ because the property of being an SP-group is quotient closed. By the induction hypothesis, G/A is nilpotent-by-finite. Therefore, G is abelian-by- (nilpotent-by-finite). We wish to show G is nilpotent-by-finite, so no harm is done in coming down to a subgroup of finite index in G — that is, we assume G is abelian-by-nilpotent. Therefore G/A is nilpotent and finitely generated, from which it follows ([9], p.132) that G/A is polycyclic.

We now invoke the theorem of P. Hall that finitely generated abelian-by-polycyclic groups satisfy *max-n*.

If G were not nilpotent-by-finite then there would exist $B \lhd G$ maximal subject to G/B not being nilpotent-by-finite, but every proper quotient of G/B is

nilpotent-by-finite. However, G/B is finitely generated, soluble, and an SP-group, so it suffices to prove that G/B is nilpotent-by-finite; equivalently we may assume hereafter that every proper quotient of G is nilpotent-by-finite. Now G is soluble and minimax (by Lemma [2.2]) so G has finite rank. This follows from the fact that polycyclic and Černikov groups have finite rank, which is an extension-closed property.

Let $\rho(A)$ denote the torsion subgroup of A. Now $\rho(A)$ is finite because G has finite rank and A is abelian. Therefore $C_G(\rho(A))$ is of finite index in G, so no harm is done if we come down to $C_G(\rho(A))$ — assume that $\rho(A) \subset \zeta(G)$. As the torsion elements are all central, clearly G is nilpotent-by-finite if and only if $G/\rho(A)$ is, too.

Conveniently, we need only condider the case that the subgroup A is torsion-free. Meanwhile G/A is polycyclic so by Lemma [3.3] we may convolute the finite pieces to the top and consider G/A as poly(C_∞) -by-finite. Once again come down to a subgroup of finite index to assume G/A is torsion-free and nilpotent, as well as finitely-generated. Therefore by Lemma [3.2], there is a set of elements $S = \{s_1, \ldots s_r\}$ of G such that

$$A = G_0 \leq \langle G_0, s_1 \rangle = G_1 \leq \cdots \leq \langle G_{r-1}, s_r \rangle = G_r = G.$$

is a central series with infinite cyclic factors. We interrupt our induction on ℓ to proceed by induction on r.

If $r = 1$, then $G_1 = G = \langle A, s_1 \rangle$. By lemma [3.4], we have that $\zeta(\langle A, s_1^{\ell_1} \rangle)$ is non-trivial for some $\ell_1 > 0$. If we set $D = A \bigcap \zeta(\langle A, s_1^{\ell_1} \rangle)$, then D is a non-trivial normal subgroup of G, so G/D is nilpotent-by-finite; moreover D centralizes $\langle A, s_1^{\ell_1} \rangle$ which is of finite index in G. Therefore, G is indeed nilpotent-by-finite.

It suffices now to prove the result for $r = d$ if the result holds for all $r < d$.

Now G_{d-1} is nilpotent-by-finite and $G_d = G = \langle G_{d-1}, s_d \rangle$. For some $\ell_1 > 0$ we have that $H = \langle A, G_{d-1}^{\ell_1} \rangle$ is a normal, nilpotent subgroup of $\langle H, s_d \rangle$. Let $M = \langle H, s_d \rangle$ and $Y = A \bigcap \zeta(H)$, then Y is normal in M. Moreover, considering the Hirsch length we see that M is of finite index in G. Using lemma [3.4] we have, for some $\ell_2 > 0$, $D = Y \bigcap \zeta(\langle Y, s_d^{\ell_2} \rangle)$ is a non-trivial subgroup of G which centralizes $\langle H, s_d^{\ell_2} \rangle$.

Now $P = \langle H, s_d^{\ell_2} \rangle$ is of finite index in G, so it contains a subgroup N of finite index in G and normal in G:

This follows from the fact that $P \leq N_G(P)$ implies that $|G : N_G(P)|$ is finite. Therefore P has a finite number of conjugates in G, say P_1, ..., P_n, (where each P_i is of finite index in G.) Using $|G : A \cap B| \leq |G : A||G : B|$ iteratively, we find that $N = \bigcap_{i=1}^{n} P_i$ is a normal subgroup of G of finite index.



Finally $D \lhd N$, so N/D is nilpotent-by-finite. In fact D is central, so N must in fact be nilpotent-by-finite; N is of finite index in G, though, so we have finally that G is nilpotent-by-finite, as desired.

Therefore if G is a finitely-generated soluble group and $G \in SP(\{u\}, \{v\})$ for words u and v, then G is nilpotent-by-finite. Q.E.D.

REFERENCES

[1] E. Hecke, "Lectures in the theory of algebraic numbers", Springer-Verlag, New York, 1981.

[2] P.H. Kropholler, *On finitely generated soluble groups with no large wreath product sections*, Proc. London Math. Soc. **(3)** 49 (1984), 155-169.

[3] A.H. Rhemtulla, *Groups with many elliptic subgroups*, to appear

[4] A.H. Rhemtulla and A.R. Weiss, *Groups with permutable subgroup products*, "Group theory: proceedings of the Singapore Group Theory Conference " (K.N. Cheng, ed.) de Gruyter, Berlin, 1989.

[5] A.H. Rhemtulla and J.S. Wilson, *Elliptically embedded subgroups of polycyclic groups*, Proc. Amer. Math. Soc. **(102)** 2 (1988) 230-234.

[6] A.H. Rhemtulla and J.S. Wilson, *On elliptically embedded subgroups of soluble groups*, Canad. J. Math. **(XXXIX)** 4 (1987), 956-968.

[7] D.J.S. Robinson, *On soluble minimax groups*, Math. Z. **(101)** (1967), 13-40.

[8] D.J.S. Robinson, "Finiteness conditions and generalized soluble groups, Parts I and II," Springer-Verlag, Berlin, 1972.

[9] D.J.S. Robinson, "A course in the theory of groups," Springer-Verlag, New-York, 1982.

[10] D. Segal, "Polycyclic groups," Cambridge University Press, Cambridge, 1983.